

*Topdressing and organic matter management for cool season golf greens*




Roch Gaussoin  
University of Nebraska-Lincoln  
[rgaussoin1@unl.edu](mailto:rgaussoin1@unl.edu)  
[@rockinsince57](https://twitter.com/rockinsince57)



Winter Webinar Series  
2023

1



**Download presentation**

2

Outline

- Historical perspective
  - Greens Construction
  - New Management Paradigm
    - Firm and Fast
    - Organic Matter Accumulation
- Fine tuning
  - Topdressing
  - Cultivation


3



4




5



6

Closer cut mowers



As low as 0.25"

7

In 1932, a fruit farmer, Orton Englehardt, invented the impact sprinkler.



8



9




10

### USGA Method of Putting Green Construction


- Original Specifications in 1960
  - Since then, this method has been regularly researched, improved and amended
- Other methods
  - California Style (1990)
  - Purr-wick (1966)
  - Dutch Green (1960-70; primarily the Netherlands)
  - Native soil or push-up greens

11


### Three Tier (USGA) System




12




13




14



15



16



17

### Root Zone Properties

Before 2004

**USGA  $K_{sat}$  guidelines**

Normal:	6-12 inches per hour
Accelerated:	12-24 inches per hour

**Account for substantial climatic differences**

Normal:	temperate to dry climates
Accelerated:	high rain subtropical and tropical climates or regions with frequent dust storms

18

Physical properties of sand-based root zones over time  
1996-2005  
University of Nebraska-Lincoln

19

**Objectives**

- Develop a better understanding of the impact of grow-in procedures on putting green establishment and performance.
- Investigate temporal changes in the soil physical properties of USGA putting greens.

20


**Materials and Methods**

- Field experiment initiated in 1997
- Greens constructed every year for four years
- Two rootzone mixtures
  - 80:20 Sand:Peat (v:v)
  - 80:15:5 Sand:Peat:Soil (v:v:v)
- Two establishment treatments
  - Accelerated
  - Controlled


21




22




23




24




25




26




27




28




29



30



31



32

### Formation of Mat

- Formation of mat layer increased approximately 0.25" (0.65 cm) annually (following establishment year).
- No visible layering, only a transition is evident between mat and original rootzone.
- Topdressing program
  - Light, Frequent
    - every 10-14 days (depending on growth) and combined with verticutting
  - Heavy, Infrequent
    - 2x annually (spring/fall) and combined with core aeration


33

### Annual organic matter accumulation in a sand/peat green

Year	1	2	3
	0.65%	3.0%	6.0%

**USGA spec. green constructed with 20% (by volume) organic matter**

34




35

### Materials and Methods

- 2004 rootzone samples taken below mat layer from each soil treatment and sent to Hummel labs for Quality Control Test (24 total samples)
- Tested against original quality control test (z-score).

36




37


### Change in Rootzone Particle Size Distribution

- All rootzones tested in 2004 showed increased proportion of fine sand (0.15 – 0.25 mm) with decreased proportion of gravel (> 2.0 mm) and very coarse sand (2.0 – 1.0 mm).

38



39



40

### Root Zone: Mat vs. Original

(samples taken July 15, 2004)

- pH: Mat < Original
- Mat > Original: CEC, OM, microbes and all nutrients

41

### Conclusions


- Based on *in situ* green testing  $K_{SAT}$  decreased, and surface moisture increased, over time due to organic matter accumulation above the original rootzone and increased fine sand content originating from topdressing sand
- Organic matter did result in positive agronomic change: pH, CEC, nutrient holding capacity, microbial stability and amount

The photograph shows a cylindrical soil core sample with a distinct dark layer at the top, representing organic matter accumulation. A yellow measuring tape is placed vertically to the right of the core, showing a height of approximately 10 inches.

42

### Want to know more?

- Gaussoin, R., R. Shearman, L. Wit, T. McClellan, and J. Lewis. 2007. Soil physical and chemical characteristics of aging golf greens. *Golf Course Manage.* 75(1):p. 161-165.



Soil physical and chemical characteristics of aging golf greens

Researcher's studied the changes in creeping bentgrass greens over an eight-year period.

43

### Research Need (2004)

- Comprehensive evaluation of sand quantity, particle size, sampling protocol and cultivation methods

44

### OM accumulates as sand greens age



1 2 3 4

8 7 6 5

Green Age (years)

45

### 8.5 year-old turf



Mat Layer

Root Zone

Sand

Sand + 20% Sphagnum

46

### Practices to change thatch into mat include topdressing and ...



47

### ... cultivation.



48



### Organic Matter Management Study

#### Objectives

1. Determine if conventional hollow tine is more effective than solid tine aerification at managing organic matter accumulation
2. Determine if venting methods are effective at managing OM accumulation

49

### Treatments

Tine Treatment	Venting Treatment
None	None
2X Hollow tine	PlanetAir
2x Solid tine	Hydroject
	Bayonet tine
	Needle tine

50

### Treatments

Tine Treatment	Venting Treatment
None	None
2X Hollow tine	PlanetAir
2x Solid tine	Hydroject
	Bayonet tine
	Needle tine

15 Trts per Rep  
6 Reps per year  
2 different years  
= A whole lot of fun for one graduate student or 180 trts

51

All treatments received the same topdressing quantity (22 ft<sup>3</sup>/M\*) but different frequency

**Equilibrated to identify differences of the practices in question**

\*1 ft<sup>3</sup> = 100 lbs of dry sand; yd<sup>3</sup> = 2700 lbs

52

### Materials and Methods


- Green Age:
  - 12 years
  - 9 years
- Data collected:
  - OM% (pre-cultivation/monthly)
  - Single wall infiltration (monthly)

53

### OM Data Analysis Year 1

- No differences between green age except for higher % in older green
- No differences among venting methods
- No interactions with solid/hollow/none

54




55

### OM Data Analysis Year 2

- No differences between green age except for higher % in older green
- No differences among venting methods
- No interactions with solid/hollow/none
- No differences among solid/hollow/none

56



57



58

### What these data do/don't suggest

- Cultivation, when topdressing quantity was equal, was insignificant as a means to control OM
- However, a superintendent must use whatever tools they have at their disposal to ensure sand is making it into the profile and not the mower buckets


59

### Topdressing interval relative to Tine/Venting combinations (22 cu ft/M)\*

- **NONE/NONE**  
– 5-10 days
- **Solid & Hollow/NONE**  
– 7-14 days
- **Solid & Hollow/Venting**  
– 14-18 days

\*Observed and calculated based on displacement and surface area opened

60



61

## Project Objective

- National Survey
- Determine cause and effect relationship among maintenance practices and their interactions relative to surface OM accumulation

62


## 2006/07/08 Samples

- Sixteen states
  - Nebraska, South Dakota, Iowa, Wyoming, Colorado, Washington, Wisconsin, Illinois, New Jersey, Minnesota, New Mexico, Montana, Hawaii, California, Connecticut, Arkansas.
- 117 golf courses sampled
  - More than 1600 samples

63



64




65

## Is the age effect misleading?

- Sampling issue:
  - Mat depth increases as green ages resulting in more OM in the same volume soil.


66



67



68



69

### Survey Summary

- None of the variables collected, by themselves, or in combination with others, predicted OM
- Courses using >18 cubic ft\*/M of topdressing with or without “venting” had lower OM
- Of the known cultivars, no differences in OM were evident

\*1 ft<sup>3</sup> = 100 lbs of dry sand; yd<sup>3</sup> = 2700 lbs

70

### Organic Matter Concentration of Creeping Bentgrass Putting Greens in the Continental U.S. and Resident Management Impact

Charles J. Schenck\*, Roch E. Gausman, and Sarah A. Gausman†

\*USGA, 1900 N. 17th St., Suite 100, Phoenix, AZ 85016; †USGA, 1900 N. 17th St., Suite 100, Phoenix, AZ 85016

**S**ince 1990, USGA has been monitoring OM concentrations in creeping bentgrass putting greens across the continental U.S. This data has been used to identify areas with high OM concentrations, which can lead to increased water infiltration, increased dry spots, and increased risk of disease. This data has also been used to identify areas with low OM concentrations, which can lead to increased erosion and increased risk of disease. This data has also been used to identify areas with high OM concentrations, which can lead to increased water infiltration, increased dry spots, and increased risk of disease. This data has also been used to identify areas with low OM concentrations, which can lead to increased erosion and increased risk of disease.

71

### Topdressing

Old Tom Morris (1821–1908) is thought to have discovered the benefits of topdressing accidentally when he spilled a wheelbarrow of sand on a putting green and noted how the turf thrived shortly afterward (Hurdzan, 2004).

J.B. Beard is his classic textbook "Turfgrass Science & Culture, 1973 writes: "The most important management practice for OM management is topdressing"

72

<https://www.usga.org/content/usga/home-page/course-care/regional-updates/central-region/2018/solid-tine-aeration-order-of-operations.html>

The image shows a screenshot of a USGA webpage. At the top, there is a URL: <https://www.usga.org/content/usga/home-page/course-care/regional-updates/central-region/2018/solid-tine-aeration-order-of-operations.html>. Below the URL is a navigation bar with links: HOME, NEWS, EVENTS, EDUCATION, GOLF, TOURS, SHOP. The main content area features a large image of a golf green being aerated by a machine. To the right of the image is a large QR code. Below the image, there is a text box that says: "Apply this order to your putting greens before solid tine aeration to improve operational efficiency." At the bottom, there is a small disclaimer: "This document is the property of every golf course management program because it contains information that is confidential and not to be shared with other golf courses." The page number 73 is located at the bottom left.

73

**"Advocates of solid-tine aeration report that they get the same benefits of thatch and organic matter reduction with less labor for the collection and removal of aeration cores. Whether you pull a core or use solid tines, it's all about sand volume and the ability to dilute organic matter in the rootzone. Regardless of the method, the most important factor is filling the hole with sand. It's all about dilution, and if you can do that with less of a mess and less labor, then solid-tine aeration is a viable alternative."**

From: <https://www.usga.org/content/usga/home-page/course-care/regional-updates/central-region/2018/solid-tine-aeration-order-of-operations.html>

The image contains a quote in bold black text. Below the quote, there is a line of text starting with "From:" followed by the same URL as in slide 73. The page number 74 is located at the bottom left.


74

- Please mark all that apply. In the last 5-10 years, on our greens, our facility has:*
- Increased topdressing quantity
  - Increased topdressing frequency
  - Increased hollow tine (equal or greater than 0.5") aeration
  - Increased solid tine (equal or greater than 0.5") aeration
  - Decreased hollow (equal or greater than 0.5") tine aeration
  - Decreased solid tine (equal or greater than 0.5") aeration
  - Made minimal changes in topdressing application quantity/frequency.
  - Made minimal changes in cultivation practices.
  - Increased "venting" practices.
- The image contains a list of seven bullet points. The first six points are on the left side, and the last three are on the right side. The text is in a standard black font. The page number 75 is located at the bottom left.


75



76



77



78

*What these data do/don't suggest*

- Cultivation, when topdressing quantity was equal, was insignificant in affecting OM
- Superintendents, however, must use **whatever tools** they have at their disposal to ensure sand is making it into the profile and not the mower buckets


79




80



81



82



83



84

### Dryject/Tine Trial Fall 2021

- Check
- Hollow ½" ID
- Solid ½"OD
- DryJect 1 (3x3)
- Needle
- DryJect 2 (3x2)
- Needle + Solid
- Needle + Hollow

Procore - 3" target depth on all tines except Dryject = 5"

Sampled day after treatment in 1' depth increments to 4 "

85

Treatment	% OM	
Check	4.5	a
Hollow	3.7	b
Needle	3.1	c
DryJect (3x3)	2.7	d
Needle + Hollow	2.3	d
DryJect (3x2)	2.3	d
Needle + Solid	2.3	d
Solid	2.2	d

- No differences among depths
- Dilution only
- Dryject and needle tine were least surface disruptive
- Hollow tine response was unexpected
- ***Data is preliminary***

86



87

### Spring 2023 Tine Trial

- ~~2~~<sup>26</sup> tine types/configurations
- 2 devices (ProCore and DryJect)
- Topdressing pre and post aeration
- Timing (spring/fall)
- OM by depth
- Surface and firmness using the USGA GS3 digital golf ball

Equipment and Tine Support Provided by

88


### What have we learned?

- A high-quality sand and a well-built root zone are relatively stable and will perform properly for many years.
- What changes over time is the surface...

89



90



**It matters how you manage the accumulating thatch/mat layer**

- Cultivation has a significant impact. At minimum, use practices that help incorporate sand.
- Topdressing is critical. Can use a fine sand (0.25-5 mm) to ensure enough sand will be applied during summer, in combo with a medium (< 1 mm) with more aggressive aerification (core, solid or injection). Avoid sands of < 0.15.

91

**Key is matching your growth rate to optimize topdressing + .....**

How much sand to use for topdressing?

- Generic recommendation is 20-40 ft<sup>3</sup> per 1000 sq. feet/yr (about 0.5 inch/M/yr)
  - UNL worked showed 20-24 ft<sup>3</sup> for OM management
- Varies by amount of:
  - Traffic
  - Grass species or cultivar
  - Nitrogen Applied
  - Water Applied
  - Microclimate/Location

92

**#clipvol “One bucket at a time”**

- Micah Woods, Asian Turfgrass Center
  - Asianturfgrass.com



93

**“Growth Potential”**


- Pace Turf
  - <https://www.paceturf.org/public/sand-and-growth-potential>



94

**Chapter 12** ASA Monograph (3RD Edition)  
**Characterization, Development, and Management of Organic Matter in Turfgrass Systems**

R.E. Gaussoin, Dep. of Agronomy and Horticulture, Univ. of Nebraska  
 W.L. Berndt, Dep. of Resort and Hospitality Management, Florida Gulf Coast University  
 C.A. Dockrell, Teagasc College of Amenity Horticulture, Dublin, Ireland  
 R.A. Drijber, Dep. of Agronomy and Horticulture, Univ. of Nebraska



95

**Acknowledgements**



96



Thank you and a great growing season!

Roch Gaussoin  
[rgaussoin1@unl.edu](mailto:rgaussoin1@unl.edu)  
[@rockinsince57](https://twitter.com/rockinsince57)