

## Golf Green Rootzone Organic Matter Management


Roch Gaussoin  
 University of Nebraska-Lincoln  
[rkaussoin1@unl.edu](mailto:rkaussoin1@unl.edu)  
 @rockinsince57



MEETING & TRADE SHOW  
 OCTOBER 15-17, 2019  
 BILLINGS CONVENTION CENTER  
 BILLINGS, MT




What do we want to learn today?



~~The~~ My organic matter journey.....

- USGA/EIFG Greens Study (9 years).
- People a lot brighter than me
  - "Talking Turf" GCSAA conversation.
  - Paul Rieke, USGA visit
  - Conversation with Paul Vermeulen. Director, Competitions Agronomy at PGA TOUR, former USGA Agronomist.
- Great funding/time support from USGA (initially), NE-GCSA, GCSA of SD, Peaks and Prairies GCSA, industry and a slew of GC supers.
- Road Show.



Organic matter development: **Are We the Problem?**

**Or:**  
*is organic matter % the green speed of the new millennium?*


**ASA Monograph (3<sup>RD</sup> Edition)**

**Chapter 12**  
**Characterization, Development, and Management of Organic Matter in Turfgrass Systems**

R.E. Gaussoin, Dep. of Agronomy and Horticulture, Univ. of Nebraska  
 W.L. Berndt, Dep. of Resort and Hospitality Management, Florida Gulf Coast University  
 C.A. Dockrell, Teagasc College of Amenity Horticulture, Dublin, Ireland  
 R.A. Drijber, Dep. of Agronomy and Horticulture, Univ. of Nebraska

How does organic matter accumulate?

- Organic matter; defined
  - dead or near dead plant residue which accumulates in the grass ecosystem



How does organic matter accumulate?

- As grasses mature there is a continual senescence of non or limited function parts (roots, shoots and leaves)
- Senescence also happens when damage or injury occurs

### How much OM is produced annually?

- Roots = 3500 lbs/acre
- Leaves = 2000 lbs/acre
- Other = 2500 lbs/acre
  
- TOTAL = **8000 lbs/acre 8967 kg/ha**

**Fairway height blue/rye estimated annual production**

### Factors influencing rootzone (P)OM accumulation

- Mowing
  - increase height=increase rooting
- Irrigation
  - root growth restricted in waterlogged soils
- Cultivation
  - increase or decrease
- Fertility
  - increase or decrease
- Stress


### Where does organic matter accumulate?




- Above ground
  - Thatch/Mat
  - clipping residue
    - relatively short term
    - "pseudo" thatch
- Below ground
  - rootzone
  - rhizosphere



### Importance of (P)OM in the rhizosphere



- deposition of particulate OM
- microbial niches
- nutrient uptake
- pathogen competition





### Thatch


A loose, intermingled, organic, layer of dead and living shoots, stems, and roots that develops between the zone of green vegetation and the soil






### Benefits of "Moderate" Thatch

- Improved resilience and cushion
- improved wear tolerance
- insulate soil/crown to temperature extremes



### Problems with excessive thatch

- Footprinting



### Problems with excessive thatch


- Scalping

### Problems with excessive thatch

- LDS




### 2010 – Wet Year – Low Organic Matter Putting Green




### Doug Soldat's work at UW

- Next 3 slides
- Wetting agents, year differences and low and high OM greens

### 2010 – Wet Year – High Organic Matter Putting Green



### 2009 – Wet Year – Low Organic Matter Putting Green




### Problems with excessive thatch

- Reduced Stress Tolerance




### Problems with excessive thatch

- Overseeding Failure



Because of inherent ambiguity in terminology and sampling techniques, the term **"thatch-mat"** has appeared frequently since the late 2000's (McCarty et al., 2007; Barton et al., 2009; Fu et al., 2009).



### Mat

Thatch that has been intermixed with mineral (soil) matter

and yet one more definition.....

SOM- Soil Organic Matter



### Physical And Chemical Characteristics Of Aging Golf Greens


Roch Gaussoin, PhD  
Jason Lewis  
Ty McClellan  
Chas Schmid  
Bob Shearman, PhD





### Treatments

- rootzone Mix
  - 80:20 (sand/peat)
  - 80:15:5 (sand/peat/soil)
- Grow-In Procedure
  - Accelerated
  - Controlled



### Project Schedule (Phase I)

1996	1997	1998	1999	2000
Greens construction (one set per year)				
Seeding				
Data collection on soil physical, chemical, and microbial characteristics influenced by rootzone materials and grow-in procedures.				



### Project Schedule (Phase II)

2002	2003	2004	2005
Data collection on soil physical and chemical characteristics as influenced by age, rootzone materials and grow-in procedures.			



## Materials and Methods

9 yr old green

10 yr old green

12 yr old green

13 yr old green


As of 2009

**Large Pores**

Significant decrease in macro-porosity over time


**Small Pores**


Significant increase in micro-porosity over time



## Formation of Mat

- Formation of mat layer increased approximately 0.65 cm annually (following establishment year).
- No visible layering, only a transition is evident between mat and original rootzone.
- Topdressing program
  - Light, Frequent
    - every 10-14 days (depending on growth) and combined with verticutting
  - Heavy, Infrequent
    - 2x annually (spring/fall) and combined with core aeration







- 2004 USGA research committee site visit
- original rootzone
- mat development

### Change in Rootzone Particle Size Distribution

- All rootzones tested in 2004 showed increased proportion of fine sand (0.15 – 0.25 mm) with decreased proportion of gravel (> 2.0 mm) and very coarse sand (2.0 – 1.0 mm).
- 5 of 8 rootzones were significant (z-score) for increased fine sand content.

### Materials and Methods

- 2004 rootzone samples taken below mat layer from each soil treatment and sent to Hummel labs for Quality Control Test (24 total samples) & tested against original quality control test (z-score).
- Other analysis also completed




### Conclusions

- The  $K_{SAT}$  decrease over time *may* be due to organic matter accumulation above and in the original rootzone and/or the increased fine sand content originating from topdressing sand

## Root Zone: Mat vs. Original

- pH:
  - Mat < Original for all USGA and California Greens.
- CEC, OM, and all Nutrients tested:
  - Mat > Original for all USGA and California Greens.



## Want to know more?


- Gaussoin, R., R. Shearman, L. Wit, T. McClellan, and J. Lewis. 2007. Soil physical and chemical characteristics of aging golf greens. *Golf Course Manage.* 75(1):p. 161-165.

all slides with the USGA logo in the corner are courtesy of Paul Vermeulen. Director, Competitions Agronomy at PGA TOUR, former USGA Agronomist


## Why is high OM considered to be “bad”?

- Loss of infiltration
- Decreased aeration
- Traps “toxic” gases
- Are these concerns real or imagined?
- Why the confusion?

## Organic Matter Recommendations




- Range
  - ✓ 1.5 – 2.5% between 0.25 to 1-inches
  - ✓ 8 – 15%
- Recommendations for almost every point in between



There is no "magic" number

- ### Analysis Methods
- Many exist, but the most relevant is "combustion" or "loss on ignition"
  - The sample represents both dead and *living* organic matter
    - Food for thought.....




### OM Testing

- Know how your sample was taken and compare notes with others that use the same protocol
- Take annual tests to determine long-term trend
  - Same time of year
  - Same location and green
- Correlate your test results with turf quality and performance during stressful environmental conditions to determine need for changes in management program
- Threshold/critical levels likely vary across the United States and from course to course

### How do you get rid of OM?

- **Decomposition (microbial)**
  - Increase surface area and aeration
  - Inoculation (???)
- **Removal**
  - Power raking, verticutting, dethatching, core aerification
- **Dilution**
  - Topdressing

### Seasonal Variation (One Location)



### Organic Matter Degradation Study

### How do you get rid of OM?

?????

### Treatments

- Aerator (Granular)
- Aerator (Liquid)
- EXP072
- EXP074
- EXP076
- Carbo-Plex
- Bio-Blend
- Carbo-Plex + Bio-Blend
- Bio-Groundskeeper (Granular)
- Bio-Groundskeeper (Liquid)
- Thatch X
- EXPO70M
- EcoChem Lawn Thatch Reducer
- Untreated Control

### Locations

- **KY Bluegrass Tee Box**
  - Native Soil
- **Bentgrass Green**
  - California Green
- **KY Bluegrass Sports Turf**
  - Sand based
  - 2.5" Mowing

### Data Collection

- **Infiltration**
- **Penotrometer**
- **Thatch**
- **Organic Matter**
  - Thatch, 0-3", 3-6"

### Plot Set-Up 14 Treatments, 3 Reps




### Analysis Summary

Source	Thatch (mm)	Clegg (g)	Infiltrat (in/hr)	Thatch (OM %)	0-3" (OM %)	3-6" (OM %)
Site	**	*	**	**	**	**
Trtmnt	NS	NS	NS	NS	NS	NS
Site X Trtmnt	NS	(0.10)	NS	NS	*	NS

### Each Plot Aerified before Application of products



### 0-3" OM %



### Summary

- No product increased degradation of OM





### How effective is removal?

- Surface disruptive, short and long term
- Core aeration is the most widespread practice recommended for OM management

Tine Size and Surface Area Chart

Tine Size (in.)	Spacing (in.)	Holes/ft <sup>2</sup>	Surface Area of One Tine	Percent Surface Area Affected
1/4	1.25 <sup>2</sup>	100	0.049	3.4%
1/4	2.5 <sup>2</sup>	25	0.049	0.9%
1/2	1.25 <sup>2</sup>	100	0.196	13.6%
1/2	2.5 <sup>2</sup>	25	0.196	3.4%
5/8	2.5 <sup>2</sup>	25	3.07	5.3%





## Sampling Procedures

### Years 1, 2, & 3

- At least 3 different greens per golf course sampled
- Soil samples taken from 3 different area per green
- Samples are evaluated for OM levels using LOI
- GPS location

- *Management & Site survey*

## Organic Matter Management

- Funded by:
  - USGA (2006)
  - Nebraska Golf Course Superintendents Assoc. (2007-2009)
  - Golf Course Superintendents Assoc. of South Dakota (2006-2009)
  - Peaks & Prairies GCSA (2007-2009)


## Sampling Methods

- Samples from at least 3 greens per course
  - (1) Problematic, (1) Non-problematic plus rebuilt or varied age/management
- 3 samples from each green
- Samples taken with  $\frac{3}{4}$  inch soil probe



## Project Objective




- National Survey
  - Determine cause and effect relationship among maintenance practices and their interactions relative to surface OM accumulation










### 2006/07/08 Samples


- **Sixteen states**
  - Nebraska, South Dakota, Iowa, Wyoming, Colorado, Washington, Wisconsin, Illinois, New Jersey, Minnesota, New Mexico, Montana, Hawaii, California, Connecticut, Arkansas.
- **117 golf courses sampled**
  - More than 1600 samples



### Is the age effect misleading?

- **Sampling issue:**
  - Mat depth increases as green ages resulting in more OM in the same volume soil.






### Survey Summary



- None of the variables collected, by themselves, or in combination with others, predicted OM
- Courses using >18 cubic ft\*/M of topdressing with or without “venting” consistently had the lowest OM
- Of the known cultivars, no differences in OM were evident


\*1 ft<sup>3</sup> = 100 lbs of dry sand; yd<sup>3</sup> = 2700 lbs



### Onsite NTEP Cultivar Evaluation


1 L-93	10 SR 1020
2 Putter	11 SR 1119
3 Cato	12 Viper
4 Crenshaw	13 Century
5 LCB-103	14 Imperial
6 Penncross	15 Penn A-1
7 Backspin	16 Penn A-4
8 Trueline	17 Penn G-6
9 Providence	18 Penn G-1





“New ultra-dense varieties of bentgrass and bermudagrass are especially susceptible to excessive organic matter accumulation due to high shoot density and the ability to grow a deep, dense root system in sand-based rootzone material. Working topdressing into a tight canopy of turf is a challenge, and much of the sand can be removed with the clippings.”

Vavrek, 2006




Topdressing and the new bents  
Easy or hard???

New bents = denser and more upright

### Pulling cores or poking holes?


- In 2005, 45 of 141 courses surveyed\* planned no core removal
  - Many had not pulled cores for 2-20 years
- 75 planned to pull cores
  - 35 of the 75 had no agronomic reason to pull a core (based on USGA Agronomist evaluation) while 40 “needed” to pull cores
- With current and evolving cultivation/topdressing/rootzone technologies is pulling a core always necessary?
- Finally, is core cultivation an effective way to manage OM ?

\* Mid-continent region USGA



### Let's take a quick look at that...

- Study initiated 22 June 2005; concluded 11 Nov. 2005
- 2 USGA Putting Greens
  - Constructed in 1997 and 2000
- Sample every 2 weeks for 20 weeks
- 2 depths of interest
  - 3 & 6”



		Time after core creation (weeks)								
		2	4	6	8	10	12	14	16	
Putting Green	Depth	Organic Matter Reduction (%)								
	9 yr	3"	79	73	70	71	69	66	67	66
		6"	71	64	60	63	59	56	57	56
5 yr	3"	73	66	61	62	62	62	61	59	
	6"	66	56	51	51	52	52	51	49	

## Organic Matter Management Study

**Objectives**

1. Determine if conventional hollow tine is more effective than solid tine aerification at managing organic matter accumulation



## Organic Matter Management Study

**Objectives**

1. Determine if conventional hollow tine is more effective than solid tine aerification at managing organic matter accumulation
2. Determine if venting (less invasive cultivation) methods are effective at managing OM accumulation

All treatments received the same topdressing quantity (22 ft<sup>3</sup>/M) but different frequency

## Treatments

Tine Treatment	Venting Treatment
None	None
2X Hollow tine	PlanetAir
2x Solid tine	Hydroject
	Bayonet tine
	Needle tine


15 Trts per Rep  
6 Reps per year  
2 different years  
= A whole lot of fun for one graduate student or 180 trts

## Regional Survey

- Annual topdressing rate for 10 well maintained courses in each region
  - Stable maintenance program during past 5 growing seasons
  - Same general turf conditions
- Surveys are not research!
  - Fertility
  - Traffic
  - Green construction
  - Etc.

### Survey Anticipated Results

- **Cool-season** - gradual increase in annual topdressing rate from north to south and east to west corresponding to length of growing season
- **Warm-season** - changes in annual topdressing rate corresponding to overseeding practices, length of growing season, annual rainfall
- Surveys are not research!
  - ✓ fertility
  - ✓ traffic
  - ✓ green construction
  - ✓ etc.




### Materials and Methods

- **Green Age:**
  - 12 years
  - 9 years
- **Data collected:**
  - OM% (pre-cultivation/monthly)
  - Single wall infiltration (monthly)


### Survey Review

- Regional differences were found, however not exactly as predicted
- Some differences did not appear to have a basis in agronomy or climatic conditions
- Survey results generate multiple questions regarding current industry practices and suggest need to conduct further research in the area of organic matter management
- It is premature to conclude that the survey provides guidance for establishing sound topdressing recommendations due to the broad range of the data and the very high standard deviation found in most regions



All treatments received the same topdressing quantity (22 ft<sup>3</sup>/M) but different frequency


**Equilibrated to identify differences of the practices in question**



### OM Data Analysis Year 1

- No differences between green age except for higher % in older green

Effect of Aerification on OM



### OM Data Analysis Year 1

- No differences between green age except for higher % in older green
- No differences among venting methods

### OM Data Analysis Year 2

- No differences between green age except for higher % in older green

### OM Data Analysis Year 1

- No differences between green age except for higher % in older green
- No differences among venting methods
- No interactions with solid/hollow/none

### OM Data Analysis Year 2

- No differences between green age except for higher % in older green
- No differences among venting methods



### OM Data Analysis Year 2

- No differences between green age except for higher % in older green
- No differences among venting methods
- No interactions with solid/hollow/none

### What these data do/don't suggest

- Topdressing is the most consistent and repeatable factor in OM management
- Cultivation was insignificant as a means to control OM
- However, a superintendent must use whatever tools they have at their disposal to insure sand is making it into the profile and not the mower buckets


### OM Data Analysis Year 2

- No differences between green age except for higher % in older green
- No differences among venting methods
- No interactions with solid/hollow/none
- No differences among solid/hollow/none

### Topdressing interval relative to t tine/vent combinations (22 cu ft/M)\*

- NONE/NONE**
  - 5-10 days
- Solid & Hollow/NONE**
  - 7-14 days
- Solid & Hollow/LIC**
  - 14-18 days

\*Observed and calculated based on displacement and surface area opened



### GreenKeeper Survey

*cool season only, mark all that apply*


**In the last 5-10 years, on our greens, our facility has:**

- Increased topdressing quantity.
- Increased topdressing frequency.
- Increased hollow (equal or greater than 0.5") tine aeration.
- Increased solid tine (equal or greater than 0.5") aeration.
- Decreased hollow (equal or greater than 0.5") tine aeration.
- Decreased solid tine (equal or greater than 0.5") aeration.
- Made minimal changes in topdressing application quantity/frequency.
- Made minimal changes in cultivation practices.
- Increased "venting" practices.

<https://www.usga.org/course-care/green-section-record/57/19/is-solid-tine-aeration-right-for-your-greens-.html>

### Is Solid-Tine Aeration Right For Your Greens?


November 20, 2018  
By Rob Stone, Regional Director, West Region



Aerating greens with solid tines is becoming more popular, which helps to speed up the aeration process while still filling the holes with sand for blade health and organic matter.



*“Advocates of solid-tine aeration report that they get the same benefits of thatch and organic matter reduction with less labor for the collection and removal of aeration cores. Whether you pull a core or use solid tines, it’s all about sand volume and the ability to dilute organic matter in the rootzone. Regardless of the method, the most important factor is filling the hole with sand. It’s all about dilution, and if you can do that with less of a mess and less labor, then solid-tine aeration is a viable alternative.”*




## Topdressing

Old Tom Morris (1821–1908) is thought to have discovered the benefits of topdressing accidentally when he spilled a wheelbarrow of sand on a putting green and noted how the turf thrived shortly afterward (Hurdzan, 2004).



J.B. Beard is his classic textbook "Turfgrass Science & Culture, 1973" writes:  
**"The most important management practice for OM management is topdressing"**



What is the "best" way to get sand into the profile?

## Layering

- Aeration alone not that effective
- Must topdress to dilute OM (change its pore size distribution) and use deficit irrigation



## Layering

- Water retention is non-uniform
- Thatch/mat layers can store twice as much water than the root zone




NOT a function of drainage

Rather it is the difference in pore size distribution among layers

**"the solution to pollution is dilution"**











**GOLF COURSE INDUSTRY**  
THE VOICE FOR TODAY'S SUPERINTENDENT

sponsored by  

Top Stories • Hot Topics • Media • Special Reports • Magazine • Subscriptions

Golf Course Industry Magazine April 2018


**True Grit**  
Feature: Special Report

In the first of a two-part series, golf managers offer their methodologies for creating greens and providing top-notch playing surfaces.

April 12, 2018

<https://www.golfcourseindustry.com/article/greens-golf-topdressing-agronomics/>

- ### How do you get rid of OM?
- Decomposition (microbial)
    - Increase surface area and aeration
    - Inoculation
  - Removal
    - Power raking, dethatching, core aeration
  - Dilution
    - Topdressing



**Next steps.....**

### When do you topdress?

	All Year	Warmer Months	Cooler Months
Northeast	57	29	14
Central	32	52	16
Southeast	55	43	2
West	45	54	1


Data presented as percentages

<https://www.golfcourseindustry.com/article/greens-golf-topdressing-agronomics/>

**Preferred particle size (mm's):**

	VC 1-2	C 0.5-1	M 0.25-0.5	F 0.15-0.5	VF 0.05-0.15	Silt/ Clay
Northeast	9	13	55	19	4	0
Central	2	11	51	35	1	0
Southeast	0	4	57	35	4	0
West	2	12	49	32	2	2

Data presented as percentages  
<https://www.golfcourseindustry.com/article/greens-golf-topdressing-agronomics/>



**Topdressing Sand:  
Sorting Out What Matters**

Nebraska Turf Conference  
 Wednesday, January 9, 2019  
 9:00 – 9:45 a.m.

James A. Murphy, Ph.D.  
 Extension Specialist in Turfgrass Management

**Frequency of Heavy Topdressing (per/yr):**

	1X	2X	3X	>3X	+ Light TD?
Northeast	17	45	25	11	85
Central	28	50	11	11	86
Southeast	18	41	21	20	86
West	19	61	12	7	86

Data presented as percentages  
<https://www.golfcourseindustry.com/article/greens-golf-topdressing-agronomics/>

**Sand Particle Size**

Particle	Diameter (mm)	Sieve Mesh #
Fine Gravel	2 – 3.4	10 – 6
V. Coarse Sand	1 – 2	18
Coarse Sand	0.5 – 1	35
Medium Sand	0.25 – 0.5	60
Fine Sand	0.15 – 0.25	100
Very Fine Sand	0.05 – 0.15	270

**Difficult to incorporate**

**Frequency of Light Topdressing (days):**

	7	14	21	28	>28	Same amount?
Northeast	10	43	15	14	18	Yes
Central	7	42	28	7	16	Yes
Southeast	32	56	6	4	2	No
West	8	41	24	13	14	Yes


Data presented as percentages  
<https://www.golfcourseindustry.com/article/greens-golf-topdressing-agronomics/>

**Particle Size Distribution for Drainage**

Particle Name	Diameter (mm)	Recommendation (by weight)
Fine Gravel	2 – 3.4	Not more than 10% total,
Very Coarse Sand	1 – 2	maximum of 3% fine gravel
Coarse Sand	0.5 – 1	Minimum of 60%
Medium Sand	0.25 – 0.5	Not more than 20%
Fine Sand	0.15 – 0.25	Not more than 5%
Very Fine Sand	0.05 – 0.15	Not more than 5%
Silt	0.002 – 0.05	Not more than 5%
Clay	< 0.002	Not more than 3%
Total Fines	very fine sand + silt + clay	Less than or equal to 10%


### Research Objectives:

- Effects of topdressing with sand lacking coarse particles (0.5-mm sand)
- Does core cultivation and backfilling holes with medium-coarse sand offset any negative effects of topdressing with sands lacking coarse particles?



### Research says, so far (3 years)...

- Topdressing improved the surface:
  - reduced the OM concentration
  - produced a drier surface
- Sand size impacts on mat layer physical properties:
  - medium-fine (>20% fine sand) increased the fineness of sand in mat layer but this did not influence infiltration or VWC
    - medium-coarse and medium-fine similar water infiltration and surface wetness
  - fine-medium sand slowed water infiltration and increased surface water retention
    - fine-medium sand substantially increased fine and very fine particles in mat layer



Sand Size	2-1 mm	1-0.5 mm	0.5-0.25 mm	0.25-0.15 mm	0.15-0.05 mm
	Very Coarse	Coarse	Medium	Fine	Very Fine
	----- % (by weight) retained -----				
Medium-coarse (1-mm)	0	30	60	10	< 1
Medium-fine (0.5-mm)	0	0	74	24	2
Fine-medium	0	4	27	48	21

### Research says, so far (3 years)...

- Core cultivation and backfilling with medium-coarse sand very effective at:
  - reduces surface wetness and OM concentration
  - reduces the amount of fine and very fine sand in the mat layer, thus offsetting the negative impact of those particles



Treatment No.	Sand Size	Factors in the Experiment			Annual Quantity of Sand Applied lbs. / 1,000-sq.-ft.
		Topdressing Rate during Growing Season lbs. / 1,000-sq.-ft.	Cultivation (twice/year; May & Oct)		
			Hollow Tine	Backfill / Topdress	
1	Medium-coarse	50	None	400	1,300
2	Medium-coarse	50	Core + Backfill	600	1,700
3	Medium-coarse	100	None	400	1,800
4	Medium-coarse	100	Core + Backfill	600	2,200
5	Medium-fine	50	None	400	1,300
6	Medium-fine	50	Core + Backfill	600	1,700
7	Medium-fine	100	None	400	1,800
8	Medium-fine	100	Core + Backfill	600	2,200
9	Fine-medium	50	None	400	1,300
10	Fine-medium	50	Core + Backfill	600	1,700
11	Fine-medium	100	None	400	1,800
12	Fine-medium	100	Core + Backfill	600	2,200
13	None	0	None	0	0
14	None	0	Core + Backfill	600	1,200

### Managing for Drier Mat Layer

#### Topdressing

- Cost and interference with play and mowing are limiting factors
- Apply as much and as often as feasible (~48 tons / acre)
- Select as coarse a sand as feasible
  - medium-fine (0.5-mm) sand with less 30% fine sand

#### Core Cultivation

- Very effective at producing a drier surface
- Needed if reducing OM is important (*allows for more sand incorporation*)\*
- Time for healing is greatest limitation (*less so for solid tines and venting*)\*


*\*Gaussoin adds*



**How much sand to use for topdressing?**


- Generic recommendation is 20-40 ft<sup>3</sup> per 1000 sq. feet/yr (about 0.5 inch/M/yr)
  - UNL worked showed 20-24 ft<sup>3</sup> for OM management
- Varies by amount of:
  - Traffic
  - Grass species or cultivar
  - Nitrogen Applied
  - Water Applied
  - Microclimate/Location

*Key is matching your growth rate to optimize topdressing + .....*



Wing Point  
GOLF & COUNTRY CLUB  
EST. 1982  
Bainbridge Island, WA

Mike Goldsberry  
Golf Course Superintendent



### Greens Organic Matter Management Tool

A Location-Based Model of Organic Matter Fate within the Sand-Based Surface Layer of a Putting Green

Ed McCoy  
Ohio State University

**Introduction**  
Managing soil organic matter (SOM) in golf course putting greens is a major agronomic challenge facing golf course superintendents. If organic matter levels become excessive, the putting surface will be soft, bumpy and prone to disease and scalping. Yet measures to control organic matter accumulation such as topdressing and core aeration are commonly disruptive and result in player dissatisfaction and reduced course revenues. This article describes a location-based simulation model of organic matter accumulation, mineralization, dilution and removal to track the fate of SOM in the sand-based surface layer of

An empirical model to predict OM fate in putting green rootzones

buckeyeturf.osu.edu

- What is it that makes greens different? Essentially, two contributing factors.
  - Sunlight = growth
  - Soil medium = growth, infiltration rate, fertilizer leaching, wilting point, firmness, cold weather tolerance, air porosity
- How do we know if each greens soil medium is different without testing for % Organic Matter?
- Why are we aerating all the greens the same if they're all different?
- What are we really trying to accomplish when we aerate our greens twice a year? Many things we all know about, but mostly it's our chance to actually incorporate sand into the profile in order to manage the ongoing accumulation of OM. We're basically using these two opportunities to balance things out.
- What's the best method for making sure we have good incorporation of sand into the aeration holes?

## Growth Potential

- #clipvol
- Pace Turf
- Micah Woods
- Bill Kreuser
- Others.....

2012 Numbers		2014 Numbers	
Green	% OM Feb, 2012	Green	% OM Feb, 2014
1	3.02	1	3.31
2	3.5	2	3.4
3	3.05	3	3.09
4	2.91	4	3.08
5	3.37	5	3.52
6	3.87	6	3.12
7	3.28	7	2.66
8	3.89	8	3.3
9	3.89	9	3.35
10	3.09	10	3.16
11	3.31	11	3.31
12	3.96	12	3.06
13	3.3	13	3.41
14	3.27	14	3.19
15	2.89	15	2.74
16	2.94	16	3.14
17	4.28	17	3.96
18	4.3	18	3.48
Putter	3.57	Putter	3.03
Chipper	4.53	Chipper	3.09

Increased sand and only solid tine implemented in 2013/14

1.) Low %OM greens received a top dressing in 3rd gear and aerated 1 time

2.) Desired %OM greens received a top dressing in 2nd gear and aerated 2 times

3.) High %OM greens received a top dressing in 1st gear and aerated 3 times

- 1st gear low was 2.5 mph
  - 38% more sand than 2nd gear
- 2nd gear low was 4 mph
  - This has been our standard gear for aeration
- 3rd gear low was 7 mph
  - 43% less sand than 2nd gear

**Pro Core 648  
3/8" solid tines**


### Organic Matter Next Steps at UNL...

- "the solution to pollution is dilution"
- Next Steps
  - Can you determine organic matter at your facility without the help of a lab?
  - Topdressing impacts on structure and fluid dynamics

Green	% OM Feb, 2012	%OM Feb, 2016
1	3.02	2.65
2	3.5	2.34
3	3.05	2.49
4	2.91	2.66
5	3.37	2.62
6	3.87	2.9
7	3.28	2.45
8	3.89	2.52
9	3.89	3.03
10	3.09	2.9
11	3.31	2.65
12	3.96	2.58
13	3.3	2.96
14	3.27	2.53
15	2.89	2.58
16	2.94	2.51
17	4.28	3.04
18	4.3	2.64
Putter	3.57	2.73
Chipper	4.53	2.56

### How is it normally done at a lab?

Loss on Ignition (LOI)





- 100-1200°C
- Sample is weighed, placed in oven, then weighed again
- OM% determined by subtraction
- Ovens are \$1200-2500

## Conclusions

- Growing medium significantly altered in 4 years.
- Infiltration rate increased.
- Lower and/or more consistent %OM throughout your greens gives you more control.
- Wilt point much higher.
- Playability is extremely consistent with golf shots being received the same throughout the course.
- We will continue to sample the greens for %OM each February and August to make informed decisions on what to do next.
- Our aeration process is simple and fast. The staff and membership are equally pleased with process and results.
- This program has become the backbone of what we do. The unit of %OM is the most important tool in our box and has allowed us to produce consistent greens that perform far better than before this program was implemented.
- Regardless of whether you pull a core or solid tine, testing each green and then knowing how much sand to use, is essential to producing consistent greens.

### Hydrogen Peroxide

- Hydrogen peroxide is a chemical compound (H<sub>2</sub>O<sub>2</sub>). In its pure form, it is a very pale blue, clear liquid, slightly more viscous than water. It is used as an **oxidizer**, bleaching agent, and antiseptic.
- Readily available & inexpensive
- Well documented to "digest" OM
- We have started testing

On course testing as an OM management tool



OM Testing

- Know how your sample was taken and compare notes with others that use the same protocol
- Take annual tests to determine long-term trend
  - Same time of year
  - Same location and green (*or all greens!*)
- Correlate your test results with turf quality and performance during stressful environmental conditions to determine need for changes in management program
- Threshold/critical levels likely vary across the United States and from course to course



Clarification/over-simplification regarding OM Management on sand based rootzones

- One size does not fit all
- The optimal % OM has not been scientifically/universally determined and may be mythical
- Cultivation is critical to increase efficiency in sand incorporation
- Solid are not different than coring tines
- The benefits of topdressing continue to be identified.

## Acknowledgements



**Nebraska Turfgrass Association**