
## Cool Season Lawn Calendar – Eastern Nebraska

A complicating factor in today's garden market is turfgrass fertilizers and preemergence weed control products are often available only as combined products. Newer (less than 20 years old) and older lawns should be managed differently for resource efficiency and environmental stewardship. This publication provides timely information for Kentucky bluegrass, tall fescue and other cool season grasses (perennial ryegrass and fine fescues.)

**Newer lawns** often struggle due to soil low in nutritional value and high weed pressure. They are best managed by making a fertilizer plus preemergence herbicide application in late April/early May followed by a second application in early June.

**Older lawns** need less fertilization to perform well and have lower weed pressure. Preemergence weed control plus fertilizer applications are optional based on need and desired outcome.

Dates	Fertilization		Cultural Practices	Pest Control	Notes
	Newer Lawn	Older Lawn			
April			Begin mowing as needed.		Mow at 3-3.5 inches all summer. Avoid removing more than 1/3 leaf height at one time.
April 25 – May 1	0.75 – 1 lbs N/1000 sq.ft.	°Optional - 0.25 lbs N/ 1000 sq.ft.	Aerate <sup>1</sup> Summer patch prevention <sup>2</sup>	Billbugs <sup>3</sup> Annual weed control – 1st application <sup>4</sup>	<sup>0</sup> If turf characteristics, such as color and density, are not ideal in early season, fertilization can be made. Otherwise, it can be skipped.
					<sup>1</sup> Use hollow tines for maximum compaction reduction.
					<sup>2</sup> When 2-inch soil depth reaches 65F temp., apply fungicides to lawns with history of infection. Repeat application 4 weeks later.
					<sup>3</sup> If lawn has a history of billbug damage. Look for adult feeding symptoms or activity in full sun near sidewalks.
					<sup>4</sup> When a 4-inch soil depth reaches 55F temp, apply fertilizer plus preemergence herbicide for annual weed control, including crabgrass and foxtail.
April 25 – May 15			Overseed <sup>5</sup>		<sup>5</sup> Core aerify before or power rake after seeding for good seed-soil contact. With new seedings, use only a preemergence products containing mesotrione or topramezone to prevent seedling death.
June 1–15	0.5 lbs N/ 1000 sq.ft.	0.5 lbs N/ 1000 sq.ft.		Preeemergence weed control – 2 <sup>nd</sup> application	



## RESOURCES

Proper Lawn Aeration – <u>https://bit.ly/CSUaeration</u> Bluegrass Billbug – <u>https://bit.ly/UMNbillbug</u> Managing Billbugs in Turfgrass – <u>https://bit.ly/PUbillbug</u> Broadleaf weed control – <u>https://go.unl.edu/NEbroadleafweed</u> Brown Patch – <u>https://byf.unl.edu/brown-patch</u> Crabgrass control – <u>https://go.unl.edu/NEcrabgrass</u> Establishing Lawns from Seed – <u>https://go.unl.edu/NEturf-seed</u> Improving Turf in Fall – <u>https://go.unl.edu/NEturf-fall</u> Irrigation audit – <u>https://go.unl.edu/NEirrigation-audit</u> Summer Patch – <u>https://bit.ly/PSUsummer-patch</u> White Grubs – <u>https://bit.ly/PUwhite-grub</u>

## AUTHORS

Sarah Browning, Horticulture Extension Educator, Nebraska Extension

John Fech, Horticulture Extension Educator, Nebraska Extension

Amanda Folck, Assistant Turfgrass Extension Educator, Department of Agronomy and Horticulture

Roch Gaussoin, Turfgrass Extension Specialist, Department of Agronomy and Horticulture