

**Ascochyta leaf blight****June 25, 2020**

Ascochyta infected turf symptoms can appear quickly and look worse than they are. Ascochyta leaf blight is considered a minor disease because injury is temporary, and rarely results in turf loss. Kentucky bluegrass is most commonly affected, but infection of other cool-season grasses, including tall fescue, is possible. Outbreaks are common throughout Nebraska.

**Symptoms.** Ascochyta may appear in isolated spots, or as large patches of straw-colored turf (similar to drought stress, but soil will likely be moist or saturated). Symptoms commonly appear following wet weather from late spring to summer, but infection may occur all season. Infections may also appear uniform and diffuse (Figure 1). Individual leaves bleach from tips, and sometimes have dark brown elliptical lesions with dark fruiting bodies called pycnidia.



**Figure 1.** Symptoms of Ascochyta leaf blight in a Kentucky bluegrass lawn (*left*) and tip dieback with pycnidia on an individual leaf blade (*right*) (Courtesy, Troy Walz, Nebraska Extension Educator).

**Management.** Disease development is most common in continuously wet turf areas, so irrigation management is critical to limit injury. Irrigate deeply and infrequently, and pre-dawn to mid-morning to limit the duration of leaf wetness. Early morning irrigation has the added benefit of knocking dew from leaves, further reducing the period of leaf wetness. Avoid evening watering. Frequent mowing, especially when turf is wet or with a dull blade increases severity. Mow when leaves are dry. Most sources indicate that fungicides are not necessary unless development is severe, and that broad-spectrum active ingredients will protect turf. We have an outbreak of Ascochyta leaf blight on a perennial ryegrass fairway on campus, and have yet to see benefit from different fungicides.

**Bottom line.** Ascochyta leaf blight may be confused with injury from drought stress, dollar spot, etc. Excessive, improperly timed irrigation and mowing increase severity. Unfortunately, knowledge of fungicide efficacy for this disease is currently limited, and the best way to prevent this disease is with sound water and mowing turf management.

**Roch Gaussoin, Professor. & Extension Turfgrass Specialist, [rgaussoin1@unl.edu](mailto:rgaussoin1@unl.edu)**